

POSITIVITY OF RICCI CURVATURE UNDER THE KÄHLER–RICCI FLOW

DAN KNOPF

1. INTRODUCTION

In analyzing a geometric evolution equation, it is of critical importance to determine which curvature conditions are preserved or attained under the flow. This is especially true for the Ricci and Kähler–Ricci flows, because of the fundamental ways in which these flows can exploit curvature conditions to reveal topological and canonical geometric properties of their underlying manifolds.

An *invariant cone* in the space of curvature operators is one that is preserved by a flow. For Ricci flow, the condition $R \geq 0$ is preserved in all dimensions, while the condition $R \leq 0$ is preserved only in real dimension two. Positive curvature operator is preserved in all dimensions [11], but positive sectional curvature is not preserved in dimensions four and above. The known counterexamples, constructed recently by Ni [18], are non-product metrics on the total space of the tangent bundles $\mathbb{R}^n \hookrightarrow TS^n \rightarrow S^n$ for $n \geq 2$. However, positive sectional curvature and positive Ricci curvature are preserved in dimension three [10]. In higher dimensions, Huisken [14] and Nishikawa [19, 20] have demonstrated the invariance of certain sets defined by curvature pinching conditions. Margerin [16] improved Huisken’s constant, thereby establishing a sharp invariant set in dimension four. Also in dimension four, H. Chen proved that 2-positivity of the curvature operator is preserved [5], while Hamilton later proved that positive isotropic curvature constitutes an invariant cone [13].

An *attractive cone* in the space of curvature operators is one that is entered asymptotically at a finite-time singularity. For example, the estimate $R \geq R_{\min}(0)$ implies that the half-space $R \geq 0$ constitutes an attractive cone in all dimensions. Attractive cones place valuable restrictions on which singularity models may appear. Important examples are the pinching estimates for 3-manifolds proved independently by Ivey [15] and Hamilton [12]. These estimates imply that any rescaled limit formed at a finite-time singularity in dimension three has nonnegative sectional curvature. In turn, this fact is of fundamental importance in Perelman’s use of κ -solutions [21, 22] in his recent spectacular progress toward the Geometrization Conjecture.

For Kähler–Ricci flow, Bando [1] and Mok [17] proved that positive bisectional curvature defines an invariant cone. Positive Ricci curvature is a more problematic condition (except of course in complex dimension one). It is not expected to be an attractive cone. Indeed, certain Kähler–Ricci solitons constructed by Feldman, Imanen, and the author [9] have Ricci curvature of mixed sign, yet are strongly

Date: February 1, 2008.

Partially supported by NSF grant DMS-0328233.

conjectured to occur as rescaled limits of singularities encountered by perturbed metrics on the Calabi manifolds $\mathbb{CP}^1 \hookrightarrow F^k \twoheadrightarrow \mathbb{CP}^{n-1}$. (See Example 2.2 of [9].)

On the other hand, it would be very useful indeed if positive Ricci curvature constituted an invariant cone for Kähler–Ricci flow. For example, parts of the proofs of some recent important results of X.-X. Chen and Tian for Kähler–Ricci flow [6, 7] only require positive Ricci curvature, but the authors impose the much stronger assumption of positive bisectional curvature to be sure that positive Ricci curvature will be preserved. (These papers also assume the existence of a Kähler–Einstein metric on the underlying manifold. See in particular Remarks 1.5 – 1.7 and Question 9.3 of [7].) While positive Ricci curvature is not expected to be preserved in high complex dimensions, there were until now promising reasons to hope that it might be preserved in complex dimension two. Unpublished work of Cao and Hamilton proves that positive orthogonal bisectional curvature $B(x, x^\perp) \geq 0$ is preserved under Kähler–Ricci flow and that its presence ensures that positive Ricci curvature is also preserved [4]. Very recent results of Phong and Sturm prove that positive Ricci curvature is preserved on a compact complex surface with the property that the sum of any two eigenvalues of its traceless curvature operator on traceless $(1, 1)$ -forms is non-negative [23]. (See also [24].)

The main purpose of this short note is to show that positive Ricci curvature is *not* preserved for arbitrary complex surfaces. We construct complete Kähler metrics of bounded curvature and non-negative Ricci curvature whose Kähler–Ricci evolutions immediately acquire Ricci curvature of mixed sign. Let L_{-1}^2 denote $\mathbb{C}^2$ blown up at the origin. (See below for an explanation of the notation.) In this paper, we will prove the following:

Theorem 1. *There exists a one-parameter family (modulo scaling) of Kähler metrics on L_{-1}^2 that are complete, of bounded curvature, and of non-negative Ricci curvature, but which immediately develop mixed Ricci curvature when evolved by the Kähler–Ricci flow.*

In fact, Theorem 1 is a special case of a more general result. For $n \geq 2$ and $k \in \mathbb{N}$, let L_{-k}^n denote the total space of the holomorphic line bundle $\mathbb{C} \hookrightarrow L_{-k}^n \twoheadrightarrow \mathbb{CP}^{n-1}$ characterized by the equation $\langle c_1, [\Sigma] \rangle = -k$, where c_1 is the first Chern class of the bundle and $\Sigma \approx \mathbb{CP}^1$ is a positively oriented generator of $H_2(\mathbb{CP}^{n-1}; \mathbb{Z})$. Since the complement of the zero-section of L_{-k}^n is biholomorphic to $(\mathbb{C}^n \setminus \{0\})/\mathbb{Z}_k$, one may construct L_{-k}^n by gluing a $\mathbb{CP}^{n-1}$ onto $(\mathbb{C}^n \setminus \{0\})/\mathbb{Z}_k$ at the origin. Notice that L_{-1}^n is simply the tautological line bundle, i.e. $\mathbb{C}^n$ blown up at the origin.

Theorem 2. *For each complex dimension $n \geq 2$ and all integers $k \in [1, n-1]$, there exists a one-parameter family (modulo scaling) of Kähler metrics on L_{-k}^n that are complete, of bounded curvature, and of non-negative Ricci curvature, but which immediately develop mixed Ricci curvature when evolved by the Kähler–Ricci flow.*

The remainder of this paper is organized as follows. Since the critical $n = 2$ case is most interesting, we begin with a detailed proof of Theorem 1. In Section 2 we provide, for the convenience of the reader, a review of $U(2)$ -invariant metrics on $\mathbb{C}^2 \setminus \{0\}$. In Section 3, we construct initial metrics on L_{-1}^2 . Then in Section 4, we finish the proof of Theorem 1. Finally, in Section 5, we indicate the straightforward generalizations by which one proves Theorem 2 for all complex dimensions $n \geq 2$.

Acknowledgement 1. *The author warmly thanks Huai-Dong Cao for encouraging him to work on this problem.*

Remark 1. *Like Ni's counterexamples for non-negative sectional curvature [18], the manifolds constructed here are noncompact. Thus it is still an open question of significant interest to find compact counterexamples for which non-negative Ricci (or sectional) curvatures are not preserved under Kähler-Ricci flow.*

2. U(2)-INVARIANT KÄHLER METRICS

In this section, we review the construction of U(2)-invariant Kähler metrics on $\mathbb{C}^2 \setminus \{0\}$. Such metrics also appear in [2, 3] and [9]. Consider $\mathbb{C}^2 \setminus \{0\}$ with complex coordinates (z_1, z_2) . Define

$$u = |z_1|^2, \quad v = |z_2|^2, \quad \text{and} \quad w = u + v.$$

Any function $P : \mathbb{C}^2 \setminus \{0\} \rightarrow \mathbb{R}$ depending only on w may be written as $P(r)$, where

$$r = \log w.$$

Anticipating that we will later abuse notation by regarding P as a function of time as well, we shall use subscripts to denote (partial) derivatives with respect to r . Define

$$(2.1) \quad \varphi(r) = P_r(r).$$

Then P is the Kähler potential for a metric g if and only if the derivatives φ and φ_r are everywhere positive. In this case, it is straightforward to compute that

$$(2.2) \quad g_{i\bar{j}} = e^{-r} \varphi \delta_{ij} + e^{-2r} (\varphi_r - \varphi) \bar{z}_i z_j$$

and

$$(2.3) \quad g^{i\bar{j}} = e^r \varphi^{-1} \delta^{ij} + (\varphi_r^{-1} - \varphi^{-1}) z^i \bar{z}^j.$$

With respect to the standard bases $(\partial/\partial z_1, \partial/\partial z_2)$ and (dz_1, dz_2) , one can therefore regard g and g^{-1} as the matrices

$$\begin{pmatrix} g_{1\bar{1}} & g_{1\bar{2}} \\ g_{2\bar{1}} & g_{2\bar{2}} \end{pmatrix} = \frac{1}{w^2} \begin{pmatrix} v\varphi + u\varphi_r & (\varphi_r - \varphi) \bar{z}_1 z_2 \\ (\varphi_r - \varphi) z_1 \bar{z}_2 & u\varphi + v\varphi_r \end{pmatrix}$$

and

$$\begin{pmatrix} g^{1\bar{1}} & g^{2\bar{1}} \\ g^{1\bar{2}} & g^{2\bar{2}} \end{pmatrix} = \frac{1}{\varphi\varphi_r} \begin{pmatrix} u\varphi + v\varphi_r & (\varphi - \varphi_r) \bar{z}_1 z_2 \\ (\varphi - \varphi_r) z_1 \bar{z}_2 & v\varphi + u\varphi_r \end{pmatrix}$$

respectively. In the standard coordinate chart, define

$$(2.4) \quad G = -\log \det g = 2r - \log \varphi - \log \varphi_r$$

and

$$(2.5) \quad \psi = G_r = 2 - \frac{\varphi_r}{\varphi} - \frac{\varphi_{rr}}{\varphi_r}.$$

Then the complex Ricci tensor $\text{Rc} = R_{i\bar{j}} dz_i d\bar{z}_j$ may be represented in the same basis as

$$(2.6) \quad \begin{pmatrix} R_{1\bar{1}} & R_{1\bar{2}} \\ R_{2\bar{1}} & R_{2\bar{2}} \end{pmatrix} = \frac{1}{w^2} \begin{pmatrix} v\psi + u\psi_r & (\psi_r - \psi) \bar{z}_1 z_2 \\ (\psi_r - \psi) z_1 \bar{z}_2 & u\psi + v\psi_r \end{pmatrix}.$$

Abusing terminology slightly, we will say V is an eigenvector of Ricci corresponding to the eigenvalue λ if $R_{ij}V^i = \lambda g_{ij}V^j$. Thus understood, the eigenvalues of Rc are

$$\lambda_1 = \frac{\psi}{w} \quad \text{and} \quad \lambda_2 = \frac{\psi_r}{w}.$$

The Levi-Civita connection of g is determined by the Christoffel symbols

$$\begin{aligned} \Gamma_{11}^1 &= \frac{\bar{z}_1}{w^2} \left[u \frac{\varphi_{rr}}{\varphi_r} + 2v \frac{\varphi_r}{\varphi} + u - 2w \right] \\ \Gamma_{12}^1 &= \frac{\bar{z}_2}{w^2} \left[u \frac{\varphi_{rr}}{\varphi_r} + (v - u) \frac{\varphi_r}{\varphi} - v \right] \\ \Gamma_{22}^1 &= \frac{z_1 \bar{z}_2^2}{w^2} \left[\frac{\varphi_{rr}}{\varphi_r} - 2 \frac{\varphi_r}{\varphi} + 1 \right] \\ \Gamma_{11}^2 &= \frac{\bar{z}_1^2 z_2}{w^2} \left[\frac{\varphi_{rr}}{\varphi_r} - 2 \frac{\varphi_r}{\varphi} + 1 \right] \\ \Gamma_{12}^2 &= \frac{\bar{z}_1}{w^2} \left[v \frac{\varphi_{rr}}{\varphi_r} + (u - v) \frac{\varphi_r}{\varphi} - u \right] \\ \Gamma_{22}^2 &= \frac{\bar{z}_2}{w^2} \left[v \frac{\varphi_{rr}}{\varphi_r} + 2u \frac{\varphi_r}{\varphi} + v - 2w \right] \end{aligned}$$

and their conjugates. The complex Riemann tensor is determined by

$$R_{i\bar{j}k\bar{\ell}} = -g_{m\bar{\ell}} \bar{\partial}_j \Gamma_{ik}^m = -\partial_i \bar{\partial}_j g_{k\bar{\ell}} + g^{p\bar{q}} \partial_i g_{k\bar{q}} \bar{\partial}_j g_{p\bar{\ell}}.$$

A straightforward computation shows that

$$\begin{aligned} R_{i\bar{j}k\bar{\ell}} &= e^{-4r} \left[-\varphi_{rrr} + 4\varphi_{rr} - 2\varphi_r + 2\varphi - 4 \frac{\varphi_r^2}{\varphi} + \frac{\varphi_{rr}^2}{\varphi_r} \right] \bar{z}_i z_j \bar{z}_k z_\ell \\ &+ e^{-3r} \left[-\varphi_{rr} + \varphi_r - \varphi + \frac{\varphi_r^2}{\varphi} \right] (\bar{z}_i z_j \delta_{k\bar{\ell}} + \bar{z}_i \delta_{jk} z_\ell + \delta_{ij} \bar{z}_k z_\ell + \delta_{i\bar{\ell}} z_j \bar{z}_k) \\ &+ e^{-2r} [-\varphi_r + \varphi] (\delta_{ij} \delta_{k\bar{\ell}} + \delta_{i\bar{\ell}} \delta_{jk}). \end{aligned}$$

To put this into a more useful form, we follow [2]. Since the metric is $U(2)$ -invariant, there is no loss of generality in evaluating the curvature at $(\zeta, 0)$, where $\zeta \neq 0$ is arbitrary. Furthermore, one may assume that the complex tangent space at $(\zeta, 0)$ is spanned by an orthonormal basis of complex vectors x, y such that $x^2 = 0$. Then one can readily compute the bisectional curvatures $B(x, x) = R(x, \bar{x}, x, \bar{x})$, $B(x, y) = R(x, \bar{x}, y, \bar{y})$, and $B(y, y) = R(y, \bar{y}, y, \bar{y})$ at $(\zeta, 0)$, obtaining

$$\begin{aligned} B(x, x)|_{(\zeta, 0)} &= \frac{1}{|\zeta|^4} \left(\frac{\varphi_{rr}^2}{\varphi_r} - \varphi_{rrr} \right) |x^1|^4 \\ B(x, y)|_{(\zeta, 0)} &= \frac{1}{|\zeta|^4} \left[\left(\frac{\varphi_{rr}^2}{\varphi_r} - \varphi_{rrr} \right) |x^1|^2 |y^1|^2 + \left(\frac{\varphi_r^2}{\varphi} - \varphi_{rr} \right) |x^1|^2 |y^2|^2 \right] \\ B(y, y)|_{(\zeta, 0)} &= \frac{1}{|\zeta|^4} \left[\left(\frac{\varphi_{rr}^2}{\varphi_r} - \varphi_{rrr} \right) |y^1|^4 + 4 \left(\frac{\varphi_r^2}{\varphi} - \varphi_{rr} \right) |y^1|^2 |y^2|^2 + 2(\varphi - \varphi_r) |y^2|^4 \right]. \end{aligned}$$

Observe in particular that the choices

$$(2.7) \quad X|_{(\zeta, 0)} = \frac{|\zeta|}{\sqrt{\varphi_r}} \frac{\partial}{\partial z_1} \quad \text{and} \quad Y|_{(\zeta, 0)} = \frac{|\zeta|}{\sqrt{\varphi}} \frac{\partial}{\partial z_2}$$

allow one to recover (2.6) in the form

$$\begin{aligned}\mathrm{Rc}(X, \bar{X})|_{(\zeta, 0)} &= B(X, X) + B(X, Y) = \frac{\psi_r}{\varphi_r} \\ \mathrm{Rc}(Y, \bar{Y})|_{(\zeta, 0)} &= B(Y, Y) + B(X, Y) = \frac{\psi}{\varphi}.\end{aligned}$$

3. AN INITIAL METRIC OF POSITIVE RICCI CURVATURE

We now want to investigate whether there exists a $U(2)$ -invariant Kähler potential P on $\mathbb{C}^2 \setminus \{0\}$ with the following global properties:

- (1) $\varphi > 0$ everywhere;
- (2) $\varphi_r > 0$ everywhere;
- (3) $\psi > 0$ everywhere;
- (4) $\psi_r \geq 0$ everywhere;
- (5) g has bounded curvature;
- (6) g extends smoothly to a complete metric as $r \rightarrow +\infty$; and
- (7) g extends smoothly to a complete metric as $r \rightarrow -\infty$.

Furthermore, we require that there exists at least one point $(\zeta, 0) \in \mathbb{C}^2 \setminus \{0\}$ such that with respect to an orthonormal basis x, y for the complex tangent space at $(\zeta, 0)$, the following local properties hold:

- I $B(x, y) < 0$; and
- II $B(x, x) = -B(x, y)$.

It may seem foolhardy to expect a single function P to satisfy all nine of these requirements. However, let $a > 0$ be a constant to be determined and consider the second-order ODE

$$(3.1) \quad [\log(\varphi\varphi_r)]_r = a.$$

Its general solution is $\sqrt{2e^{ar+b}/a+c}$, where b and c are arbitrary. Without loss of generality, one may eliminate b by scaling, yielding

$$(3.2) \quad \varphi(r) = \sqrt{\frac{2}{a}e^{ar} + c}.$$

One then has

$$(3.3) \quad \varphi\varphi_r = e^{ar},$$

so that global properties (1) and (2) are satisfied whenever $c \geq 0$. Moreover,

$$\psi = 2 - a$$

everywhere. So properties (3) and (4) are satisfied whenever $a \in (0, 2)$.

To evaluate the bisectional curvatures of the resulting metric, it suffices to consider the orthonormal basis (2.7) for the complex tangent space at $(\zeta, 0)$, where $\zeta \neq 0$ is arbitrary. One finds that

$$(3.4) \quad B(X, X) = \frac{ac}{\varphi^3} = -B(X, Y),$$

so that local properties I and II will in fact be satisfied globally if $c > 0$. The choices $a \in (0, 2)$ and $c > 0$ also ensure that

$$B(Y, Y) = \frac{2}{\varphi^3} \left[\left(\frac{2}{a} - 1 \right) e^{ar} + c \right] > B(X, X).$$

Because $\varphi(r) \rightarrow \sqrt{c} > 0$ as $r \rightarrow -\infty$ and $\varphi(r) = \mathcal{O}(e^{(a/2)r})$ as $r \rightarrow \infty$, the curvatures are globally bounded, satisfying property (5). To investigate completeness at spatial infinity, observe that (3.3) implies that $\varphi_r = \mathcal{O}(e^{ar})$ as $r \rightarrow -\infty$ and that $\varphi_r = \mathcal{O}(e^{(a/2)r})$ as $r \rightarrow \infty$. Since radial paths emanating from the origin are geodesics and the length of such a path is $\int_{-\infty}^{\infty} \sqrt{\varphi_r} dr$, it is easy to see that the metric is complete as $r \rightarrow +\infty$, satisfying global property (6).

On the other hand, the metric distance from an arbitrary point to $r = -\infty$ is finite. Our choice of $c > 0$ makes it impossible to complete the metric smoothly by gluing in a point at the origin. But Calabi's theorem [8] tells us that g will extend to a smooth Kähler metric on L^2_{-k} if φ has an asymptotic expansion near $w = 0$ of the form

$$\varphi = a_0 + a_1 w^k + a_2 w^{2k} + \mathcal{O}(w^{3k}),$$

with a_0 and a_1 both positive. (See Section 4.1 of [9] for an exposition.) The function (3.2) will have such an expansion if a is a positive integer. Fortunately for us, there is a (unique) positive integer in the interval $(0, 2)$. Taking $a = 1$ thus lets us satisfy global property (7). Notice that for each $c > 0$, the solution φ then admits the expansion

$$\varphi = \sqrt{c} + \frac{1}{\sqrt{c}} w - \frac{1}{2c^{3/2}} w^2 + \cdots.$$

In particular, $2\pi\sqrt{c}$ is the area of the $\mathbb{CP}^1$ that constitutes the zero-section of the bundle.

Remark 2. *The Kähler manifolds (L^2_{-1}, g) we have constructed are asymptotic at infinity to a Kähler cone Γ on $\mathbb{C}^2 \setminus \{0\}$ whose cone angle around the origin in each complex line is π and whose density ratio is $\text{Vol}_\Gamma(B_1)/\text{Vol}_{\mathbb{C}^2}(B_1) = 1/4$.*

4. EFFECT OF THE KÄHLER-RICCI FLOW

We now consider the Kähler-Ricci evolution $(L^2_{-1}, g(t))$, where we take $P(r, 0)$ to be the Kähler potential $P(r)$ constructed above. Again choose an arbitrary point $(\zeta, 0) \in \mathbb{C}^2 \setminus \{0\}$. Because

$$g|_{(\zeta, 0)} = \frac{1}{|\zeta|^2} \begin{pmatrix} \varphi_r & 0 \\ 0 & \varphi \end{pmatrix}$$

and

$$\text{Rc}|_{(\zeta, 0)} = \frac{1}{|\zeta|^2} \begin{pmatrix} \psi_r & 0 \\ 0 & \psi \end{pmatrix}$$

in the standard basis, the Kähler-Ricci flow

$$\frac{\partial}{\partial t} g = -\text{Rc}$$

will be satisfied if and only if φ evolves by the PDE $\varphi_t = -\psi$; to wit,

$$(4.1) \quad \varphi_t = \frac{\varphi_{rr}}{\varphi_r} + \frac{\varphi_r}{\varphi} - 2.$$

Since $\partial/\partial r$ and $\partial/\partial t$ commute, it follows that the Ricci potential ψ must evolve by the PDE

$$(4.2) \quad \psi_t = \frac{\psi_{rr}}{\varphi_r} - \frac{\varphi_{rr}\psi_r}{\varphi_r^2} + \frac{\psi_r}{\varphi} - \frac{\varphi_r\psi}{\varphi^2}.$$

Since $\psi(\cdot, 0) \equiv 1$, this reduces to

$$\left. \frac{\partial}{\partial t} \psi \right|_{t=0} = -\frac{\varphi_r}{\varphi^2}$$

at the initial time. A further computation then shows that

$$(4.3) \quad \left. \frac{\partial}{\partial t} \psi_r \right|_{t=0} = \frac{e^r}{\varphi^5} (e^r - c),$$

which is strictly negative for all $r < \log c$. Since $\psi_r(\cdot, 0) \equiv 0$, the complex Ricci tensor must acquire a negative eigenvalue $\lambda_2 < 0$ at all such points for small times $t > 0$. This completes the proof of Theorem 1.

5. THE GENERAL CASE: COMPLEX DIMENSION $n \geq 2$

If P is a $U(n)$ -invariant Kähler potential on $\mathbb{C}^n \setminus \{0\}$, formulas (2.2) for g and (2.3) for g^{-1} remain unchanged. Formula (2.4) becomes

$$G = nr - \log \varphi - \log \varphi_r,$$

whence (2.5) is replaced by

$$\psi = G_r = n - \frac{\varphi_r}{\varphi} - \frac{\varphi_{rr}}{\varphi_r}.$$

One calculates easily that

$$R_{i\bar{j}} = e^{-r} \psi \delta_{ij} + e^{-2r} (\psi_r - \psi) \bar{z}_i z_j.$$

To display the eigenvalues of Rc , one may without loss of generality fix $Z = (\zeta, 0, \dots, 0) \in \mathbb{C}^n \setminus \{0\}$ with $\zeta \neq 0$ arbitrary. Then

$$g|_Z = \frac{1}{|\zeta|^2} \begin{bmatrix} \varphi_r & & & \\ & \varphi & & \\ & & \ddots & \\ & & & \varphi \end{bmatrix}$$

and

$$\text{Rc}|_Z = \frac{1}{|\zeta|^2} \begin{bmatrix} \psi_r & & & \\ & \psi & & \\ & & \ddots & \\ & & & \psi \end{bmatrix}.$$

If φ is defined by (3.2) with $a > 0$, one finds that

$$\psi \equiv n - a,$$

hence that $\psi_r \equiv 0$. Equation (4.1) becomes

$$\varphi_t = \frac{\varphi_{rr}}{\varphi_r} + \frac{\varphi_r}{\varphi} - n,$$

so that (4.2) remains unchanged. The analog of (4.3) is

$$\left. \frac{\partial}{\partial t} \psi_r \right|_{t=0} = (n - a) \frac{e^{ar}}{\varphi^5} (e^{ar} - ac),$$

which is strictly negative for all $a \in (0, n)$ and $c > 0$ whenever $r < \frac{1}{a} \log(ac)$. If we take $a \in (0, n)$ to be an integer, then φ will have an asymptotic expansion near the origin of the form

$$\varphi = \sqrt{c} + \frac{1}{a\sqrt{c}}w^a - \frac{1}{2a^2c^{3/2}}w^{2a} + \dots.$$

Hence by Calabi's theorem, g will extend to a smooth metric on L_{-a}^n for all such a . It is easy to check that g is complete and asymptotic to a Kähler cone at infinity, hence of bounded curvature. This completes the proof of Theorem 2.

Remark 3. For each $1 \leq k \leq n-1$, the Kähler manifolds (L_{-k}^n, g) we have constructed are asymptotic at infinity to a Kähler cone Γ_k on $\mathbb{C}^n \setminus \{0\}$ whose cone angle around the origin is π and whose density ratio is $\text{Vol}_{\Gamma_k}(B_1)/\text{Vol}_{\mathbb{C}^n}(B_1) = k^{1-n}/2^n$.

REFERENCES

- [1] **Bando, Shigetoshi.** On the classification of three-dimensional compact Kaehler manifolds of nonnegative bisectional curvature. *J. Differential Geom.* **19** (1984), no. 2, 283–297.
- [2] **Cao, Huai-Dong.** Existence of gradient Kähler-Ricci solitons. *Elliptic and parabolic methods in geometry (Minneapolis, MN, 1994)*, 1–16, *A K Peters*, Wellesley, MA, 1996.
- [3] **Cao, Huai-Dong.** Limits of solutions to the Kähler-Ricci flow. *J. Differential Geom.* **45** (1997), no. 2, 257–272.
- [4] **Cao, Huai-Dong; Hamilton, Richard S.** On the preservation of positive orthogonal bisectional curvature under the Kähler-Ricci flow. (1988) Unpublished.
- [5] **Chen, Haiwen.** Pointwise $\frac{1}{4}$ -pinched 4-manifolds. *Ann. Global Anal. Geom.* **9** (1991), no. 2, 161–176.
- [6] **Chen, Xiuxiong; Tian, Gang.** Ricci flow on Kähler-Einstein surfaces. *Invent Math.* **147** (2002), no. 3, 487–544.
- [7] **Chen, Xiuxiong; Tian, Gang.** Ricci flow on Kähler-Einstein manifolds. [arXiv:math.DG/0108179](#).
- [8] **Calabi, Eugenio.** Extremal Kähler metrics. *Seminar on Differential Geometry*, 259–290, *Ann. of Math. Stud.*, **102**, *Princeton Univ. Press*, Princeton, N.J., 1982.
- [9] **Feldman, Mikhail; Ilmanen, Tom; Knopf, Dan.** Rotationally symmetric shrinking and expanding gradient Kähler-Ricci solitons. *J. Differential Geom.* **65** (2003), no. 2, 169–209.
- [10] **Hamilton, Richard S.** Three-manifolds with positive Ricci curvature. *J. Differential Geom.* **17** (1982), no. 2, 255–306.
- [11] **Hamilton, Richard S.** Four-manifolds with positive curvature operator. *J. Differential Geom.* **24** (1986), no. 2, 153–179.
- [12] **Hamilton, Richard S.** The formation of singularities in the Ricci flow. *Surveys in differential geometry*, Vol. II (Cambridge, MA, 1993), 7–136, *Internat. Press*, Cambridge, MA, 1995.
- [13] **Hamilton, Richard S.** Four-manifolds with positive isotropic curvature. *Comm. Anal. Geom.* **5** (1997), no. 1, 1–92.
- [14] **Huisken, Gerhard.** Ricci deformation of the metric on a Riemannian manifold. *J. Differential Geom.* **21** (1985), no. 1, 47–62.
- [15] **Ivey, Thomas.** Ricci solitons on compact three-manifolds. *Differential Geom. Appl.* **3** (1993), no. 4, 301–307.
- [16] **Margerin, Christophe.** A sharp characterization of the smooth 4-sphere in curvature terms. *Comm. Anal. Geom.* **6** (1998), no. 1, 21–65.
- [17] **Mok, Ngaiming.** The uniformization theorem for compact Kähler manifolds of nonnegative holomorphic bisectional curvature. *J. Differential Geom.* **27** (1988), no. 2, 179–214.
- [18] **Ni, Lei.** Ricci flow and nonnegativity of curvature. [arXiv:math.DG/0305246](#).
- [19] **Nishikawa, Seiki.** Deformation of Riemannian metrics and manifolds with bounded curvature ratios. *Geometric measure theory and the calculus of variations (Arcata, Calif., 1984)*, 343–352, *Proc. Sympos. Pure Math.*, **44**, *Amer. Math. Soc.*, Providence, RI, 1986.

- [20] **Nishikawa, Seiki.** On deformation of Riemannian metrics and manifolds with positive curvature operator. *Curvature and topology of Riemannian manifolds (Katata, 1985)*, 202–211, Lecture Notes in Math., **1201**, Springer, Berlin, 1986.
- [21] **Perelman, Grisha.** The entropy formula for the Ricci flow and its geometric applications. [arXiv:math.DG/0211159](#).
- [22] **Perelman, Grisha.** Ricci flow with surgery on three-manifolds. [arXiv:math.DG/0303109](#).
- [23] **Phong, Duong H.; Sturm, Jacob.** On the Kähler-Ricci flow on complex surfaces. [arXiv:math.DG/0407232](#).
- [24] **Phong, Duong H.; Sturm, Jacob.** On stability and the convergence of the Kähler-Ricci flow. [arXiv:math.DG/0412185](#).

UNIVERSITY OF TEXAS AT AUSTIN

E-mail address: danknopf@math.utexas.edu

URL: <http://www.ma.utexas.edu/~danknopf/>